

# Hadronic vacuum polarization contribution to muon $g-2$ using staggered fermions

Christopher Aubin  
June 18, 2012  
Project X

with  
Tom Blum, Maarten  
Golterman, & Santi Peris




FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK



# “Theory” vs. Experiment

Davies, arXiv:1001.2243



Deviation  $\sim 3$ -3.8 sigma

Fermilab experiment will start in **2016**, to reduce error from 0.5 to 0.1 ppm

$$a_{\mu}^{\text{exp}} = \left( \frac{g_{\mu} - 2}{2} \right)^{\text{exp}} = 11\,659\,208(6) \times 10^{-10}$$



**FORDHAM UNIVERSITY**  
THE JESUIT UNIVERSITY OF NEW YORK

# Outline

Muon  $g-2$  and current theory for HLO

HLO:  $O(\alpha^2)$  Contribution—Vacuum Polarization

$(g-2)^{\text{LHO}}$  from first principles:  
Lattice Gauge Theory



Lattice results for vacuum polarization

Preliminary fits for  $g-2$   
VMD vs. Padé Approximants

# Muon g-2

$$\Gamma^\mu = \gamma^\mu F_1(q^2) + \frac{i\sigma^{\mu\nu}}{2m_\mu} F_2(q^2)$$

$$a_\mu = \left( \frac{g_\mu - 2}{2} \right) = F_2(q^2 = 0)$$




+ other non-QCD terms...




**FORDHAM UNIVERSITY**  
THE JESUIT UNIVERSITY OF NEW YORK

# Hadronic contributions

Vacuum Polarization



Light by light



I'll focus on the *leading* hadronic contribution, the vacuum polarization



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

# Leading Hadronic Contribution

The  $O(\alpha^2)$  hadronic contribution,  $a_\mu^{\text{HLO}}$ , cannot be calculated in perturbation theory.

Via the Optical theorem, one can evaluate it using  $\sigma(e^+e^- \rightarrow \text{hadrons})$ .

$$a_\mu^{\text{HLO}} = \frac{\alpha^2}{3\pi^2} \int_{4m_\pi^2}^{\infty} \frac{ds}{s} K(s) R(s)$$

The kernel,  $K(s)$ , is known, and  $R(s)$  can be measured experimentally

Not a theoretical problem since 1961!




FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK



$$R(s) = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)}$$

The precision of the Standard Model prediction is limited by the experimental measurement of  $R(s)$ .



(from Davier et al, hep-ph/0208177)



# Lattice QCD

Use staggered quarks (MILC collaboration):

Large volumes  
Small quark masses  
High statistics

Asqtad staggered (reduced lattice spacing errors)  
3 lattice spacings (that we use)  
pion masses as low as 180 MeV


Future: HISQ quarks with nearly physical pion masses



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK



# Leading Hadronic Contribution



We can extract the  $O(\alpha^2)$  hadronic contribution to  $a_\mu$  from the vacuum polarization using the Euclidean space expression (Blum, 2003)

$$a_\mu^{(2)\text{had,LO}} = \left(\frac{\alpha}{\pi}\right)^2 \int_0^\infty dK^2 f(K^2) [\Pi(K^2) - \Pi(0)]$$

$f(K^2)$  diverges as  $K^2 \rightarrow 0 \Rightarrow$  dominated by low momentum region  
 $\Rightarrow$  Need large lattices to simulate these low momenta accurately



**FORDHAM UNIVERSITY**  
THE JESUIT UNIVERSITY OF NEW YORK

# Lattice Calculation of $\Pi^{\mu\nu}(q)$

Calculate the vacuum polarization using the conserved current

$$\Pi^{\mu\nu}(q) = \int d^4x e^{iq \cdot (x-y)} \langle J^\mu(x) J^\nu(y) \rangle = (q^2 g^{\mu\nu} - q^\mu q^\nu) \Pi(q^2)$$

In the continuum we have the conserved (local) EM current:

$$J^\mu = \bar{\psi} \gamma^\mu \psi$$

While on the lattice it is a point-split current:

$$J_{\mu \ x} = \frac{1}{2} \left[ \bar{\psi}_{x+a\hat{\mu}} U_{\mu x}^\dagger (1 + \gamma_\mu) \psi_x - \bar{\psi}_x U_{\mu x} (1 - \gamma_\mu) \psi_{x+a\hat{\mu}} \right]$$

Satisfies:

$$\frac{1}{a} \sum_\mu [J_{\mu \ x} - J_{\mu \ x-a\hat{\mu}}] = 0$$



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

# Lattice Calculation of $\Pi^{\mu\nu}(q)$

On the lattice this satisfies a discrete Ward Identity, so we modify the expressions above

$$Q_\mu \rightarrow \hat{Q}_\mu = \frac{2}{a} \sin \left( \frac{aQ_\mu}{2} \right)$$

The finite volume restricts the momenta:

$$Q_\mu = \frac{2\pi n_\mu}{aL_\mu}$$

SO

$$\Pi^{\mu\nu}(Q) = (\hat{Q}_\mu \hat{Q}_\nu - \hat{Q}^2 \delta_{\mu\nu}) \Pi(Q^2)$$


This provides a strong check on the simulation!



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

# Lattice Calculation of $\Pi^{\mu\nu}(q)$

To perform lattice calculation:  
Wick contract the quark fields in the current, giving two types of contractions:



We neglect second contraction *for now* (suppressed, also very noisy)

Hard to fit low-momentum region — Also most important part



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

(No lattice talk is complete without a table of numbers in unphysical units only the speaker understands)

2+1-flavor MILC “Asqtad” staggered configurations

$a$ (fm)	$am_l$	$am_s$	$\beta$	size	$m_\pi L$	# lats. avail.
$\approx 0.09^*$	0.0124	0.031	7.11	$28^3 \times 96$	5.78	531
$\approx 0.09^*$	0.0062	0.031	7.09	$28^3 \times 96$	4.14	591
$\approx 0.09$	0.00465	0.031	7.085	$32^3 \times 96$	4.10	480
$\approx 0.09^*$	0.0031	0.031	7.08	$40^3 \times 96$	4.22	945
$\approx 0.09^*$	0.00155	0.031	7.075	$64^3 \times 96$	4.80	491
$\approx \mathbf{0.09}$	0.0031	0.0031	7.045	$40^3 \times 96$	4.20	440
$\approx 0.06^\dagger$	0.0036	0.018	7.47	$48^3 \times 144$	4.50	751
$\approx 0.06^\dagger$	0.0025	0.018	7.465	$56^3 \times 144$	4.38	768
$\approx 0.06^*$	0.0018	0.018	7.46	$64^3 \times 144$	4.27	826
$\approx \mathbf{0.045}$	0.0028	0.014	7.81	$64^3 \times 192$	4.56	801

Now: 3 lattice spacings, and pion masses as low as  
~180 MeV




THE JESUIT UNIVERSITY OF NEW YORK

All (new) simulations were performed using USQCD  
Lattice resources here at Fermilab  
(the Ds and J/Psi clusters)



# All results (thus far)



# Fitting

Can't calculate the vacuum polarization at  $q^2=0$  directly

For high momentum, continuum PT works

For low momentum:

Simple polynomials?

No good beyond cubic/quartic order in  $q^2$

Vector Meson Dominance/ChPT

Padé Approximants



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

# Vector Meson Dominance

$$\Pi(Q^2) = \frac{A}{Q^2 + m_\rho^2} + B$$

$A$  is related to rho decay constant – in principle could determine both this and the rho mass from simulations, thus only  $B$  is a free parameter

For small pion masses, the two-pion state is the lightest state in this channel – can't measure rho mass (yet)!



# Rho decay

On our lightest fine ( $a=0.09$  fm) lattice, the rho can “decay”

Here, the rho mass cannot be extracted very easily, since

$$am_{\rho}^{\text{phys}} \approx 0.35 \ (\approx am_{\rho}^{\text{lat}})$$

And the two smallest non-zero spatial momenta allowed are

$$ap = \frac{2\pi}{L}, \frac{2\sqrt{2}\pi}{L}$$


For a two pion state with rho quantum numbers:

$$aE_{2\pi} = 0.2374, 0.3081 < am_{\rho}$$



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

Can be seen to affect the vacuum polarization!



This must be understood theoretically, as it clearly effects the low-energy regime – *cannot get a fully reliable result without it!*

Also tricky to use rho mass as a fixed parameter in fits – without a complete study of mixing with 2-pion state

$$\begin{pmatrix} \langle \rho | \rho \rangle & \langle \rho | 2\pi \rangle \\ \langle 2\pi | \rho \rangle & \langle 2\pi | 2\pi \rangle \end{pmatrix}$$

Need a full finite-volume analysis to disentangle the rho and 2 pion states



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK



$$\frac{\Pi(0) - \Pi(Q^2)}{Q^2} = \int_{4m_\pi^2}^{\infty} dt \frac{\rho(t)}{t(t + Q^2)} \equiv \Phi(Q^2)$$

is a *Stieltjes* function, analytic at all points not on the cut  $(-\infty, -4m_\pi^2]$

## Theorem:

Given  $P$  points  $(Q_i^2, \Phi(Q_i^2))$ , a sequence of Padé Approximants can be constructed which converge to  $\Phi(Q^2)$  on any closed, bounded region of the complex plane excluding the cut, in the limit  $P \rightarrow \infty$ .  
(Baker 1969, Barnsley 1973)



# Padé Approximants

This can be written as

$$\Pi(Q^2) = \Pi(0) - Q^2 \left( a_0 + \sum_{n=1}^{\lfloor P/2 \rfloor} \frac{a_n}{b_n + Q^2} \right)$$

$$a_{n>0} > 0, \quad b_{\lfloor P/2 \rfloor} > \cdots > b_1 > 4m_\pi^2$$

if P is even:  $a_0 = 0$

For different values of P, we fit to different Padé's

P	2	3	4	5
Padé	[0,1]	[1,1]	[1,2]	[2,2]

For comparison, we will cut off the integral for the g-2 at (1 GeV)<sup>2</sup>

Note that VMD is a [0,1] Padé, but with its pole fixed to be the vector mass, and as such **is not a valid Padé for our purposes!**



**FORDHAM UNIVERSITY**  
THE JESUIT UNIVERSITY OF NEW YORK

# Test on fine MILC lattices (pion mass = 480 MeV)

		correlated interval $0 < Q^2 \leq 0.6 \text{ GeV}^2$		uncorrelated interval $0 < Q^2 \leq 1 \text{ GeV}^2$	
PA	# parameters	$\chi^2/\text{dof}$	$10^{10} a_\mu^{\text{HLO}, Q^2 \leq 1}$	$\chi^2/\text{dof}$	$10^{10} a_\mu^{\text{HLO}, Q^2 \leq 1}$
VMD	2	5.86/3*	363(7)	4.37/18	413(8)
[0, 1]	3	11.4/8	338(6)	3.58/17	373(37)
[1, 1]	4	7.49/7	350(8)	3.36/16	424(116)
[1, 2]	5	7.49/6	350(8)	3.35/15	443(293)
[2, 2]	6	7.49/5	350(7)	3.35/14	445(432)

\* interval  $0 < Q^2 \leq 0.35 \text{ GeV}^2$


Correlated Padé's are stable – better with more parameters  
Higher poles ill-determined (does not affect g-2)

Consistent unless one compares uncorrelated VMD with the correlated fits




FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

[1,1] Padé, corr (solid) vs. uncorr (dashed)



[1,1] Padé, corr (solid) vs. VMD uncorr (dashed)



Correlated fits systematically low  
All fits have reasonable  $\chi^2$

“By eye” – no way to choose  
one fit over another

Superfine results:

$$a = 0.06 \text{ fm} \quad m_\pi = 220 \text{ MeV}$$

[1,1] Padé (corr):

$$a_\mu^{\text{HLO}, Q^2 \leq 1} = 572(41) \times 10^{10}$$


VMD (uncorr):

$$a_\mu^{\text{HLO}, Q^2 \leq 1} = 646(8) \times 10^{10}$$



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

# Difficulty



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

Note that fits can be misleading!

Unknown systematics are hidden in VMD fits

Any fits which use data primarily excluding low momentum region should be met with caution!

17% discrepancy between VMD & Padé fits

Primary Problems:

Low momentum (Large volumes/TBC's)

Statistics (AMA)

Disconnected contributions (definitely essential for ~5% unc)

Light quark masses (soon not a problem)

Chiral Extrapolation?

Soon to be irrelevant – HISQ ensembles with near-physical pion mass



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK



Note that fits can be misleading!

Unknown systematics are hidden in VMD fits

Any fits which use data primarily excluding low momentum region should be met with caution!

17% discrepancy between VMD & Padé fits

Primary Problems:

Low momentum (Large volumes/TBC's)

Statistics (AMA)

Disconnected contributions (definitely essential for ~5% unc)

Light quark masses (soon not a problem)

Non-trivial problem!

Chiral Extrapolation?

Soon to be irrelevant – HISQ ensembles with near-physical pion mass



FORDHAM UNIVERSITY  
THE JESUIT UNIVERSITY OF NEW YORK

## Conclusions

Full results still yet to come, analysis complicated by light masses...

### Immediate future:

- Better statistics using all-mode averaging (in progress)

- Thus improved fits (able to get higher Padé poles?)

- Begin simulations on HISQ ensembles with nearly physical pion mass

### Additionally:

- Need to fill in low momentum region (twisted boundary conditions)

### Longer term:

- Include disconnected diagrams

Stay tuned!



**FORDHAM UNIVERSITY**  
THE JESUIT UNIVERSITY OF NEW YORK